

A safety net for fast migrators: Interactions between gap-opening and sub-gap-opening bodies in a protoplanetary disk

Edward W. Thommes

*Canadian Institute for Theoretical Astronomy, University of Toronto, 60 St. George Street,
Toronto, Ontario M5S 3H8*

thommes@cita.utoronto.ca

ABSTRACT

Young planets interact with their parent gas disks through tidal torques. An imbalance between inner and outer torques causes bodies of mass $\gtrsim 0.1$ Earth masses ($M_{\oplus}$) to lose angular momentum and migrate inward rapidly relative to the disk; this is known as “Type I” migration. However, protoplanets that grow to gas giant mass, $O(10^2) M_{\oplus}$, open a gap in the disk and are subsequently constrained to migrate more slowly, locked into the disk’s viscous evolution in what is called “Type II” migration. In a young planetary system, both Type I and Type II bodies likely coexist; if so, differential migration ought to result in close encounters when the former originate on orbits exterior to the latter. We investigate the resulting dynamics, using two different numerical approaches: an N-body code with dissipative forces added to simulate the effect of the gas disk, and a hybrid code which combines an N-body component with a 1-dimensional viscous disk model, treating planet-disk interactions in a more self-consistent manner. In both cases, we find that sub-gap-opening bodies have a high likelihood of being resonantly captured when they encounter a gap-opening body. A giant planet thus tends to act as a barrier in a protoplanetary disk, collecting smaller protoplanets outside of its orbit. Such behavior has two important implications for giant planet formation: First, for captured protoplanets it mitigates the problem of the migration timescale becoming shorter than the growth timescale when $M_{\text{proto}} \gtrsim 1M_{\oplus}$. Secondly, it suggests one path to forming systems with multiple giant planets: Once the first has formed, it traps (or indeed accretes) the future solid core of the second in an exterior mean-motion resonance, and so on. The most critical step in giant planet formation may thus be the formation of the very first one.

Subject headings: solar system: formation — planets and satellites: formation — planetary systems: protoplanetary disks

1. Introduction

Observational evidence strongly suggests that the formation of giant planets is a ubiquitous process. To date, well over a hundred extrasolar planets have been discovered by radial velocity surveys; in at least eleven cases, multiple planets orbit a single star¹. At least three of the multi-planet systems have pairs of planets locked in mean-motion resonances (MMRs), with a 1:2 (GJ 876b,c; HD 82943b,c) or 1:3 (55 Cnc b,c) period ratio.

Migration due to planet-disk interaction is the leading candidate for the mechanism which produces such dynamical configurations, as well as producing the large fraction of “hot Jupiters”, exoplanets on orbits significantly smaller than their likely formation distance from their parent star. Indeed, the latter effect was first predicted, and investigated in detail, well before it was observed (Goldreich & Tremaine 1980; Lin & Papaloizou 1986). More recent work has expanded the focus from planet-disk interaction to “planet-planet-disk” dynamics, that is, the behavior of multiple interacting planets embedded in a disk. Again, theoretical work anticipated observation: Using hydrodynamic simulations, Kley (2000) and Bryden et al. (2000) demonstrated convergent planet migration, thus raising the possibility of resonant capture, just prior to the discovery of the first extrasolar MMR, the 1:2 resonant pair of GJ876b and c. Subsequent work (Masset & Snellgrove 2001; Lee & Peale 2002; Murray, Paskowitz, & Holman 2002; Nelson & Papaloizou 2002; Kley, Peitz, & Bryden 2004) looked in more detail on the dynamics of capture and evolution of gas giant planets in low-order MMRs.

The above work focused on the interaction of pairs of *gap-opening* planets, that is, planets which have grown large enough to lock themselves into the viscous evolution of their parent disk. For typical protoplanetary disk parameters, this implies planets of gas giant (Jovian) size, $O(10^2)$ Earth masses ($M_{\oplus}$), rather than ice giant (Neptunian) size, $O(10^1) M_{\oplus}$ planets. For initially well-separated planets, convergent migration then requires the removal of the intervening gas, thus bringing the planets to share a common gap. It is possible, however, for giant planet MMRs to be established earlier. Prior to opening a gap, disk tidal torques can cause the orbital decay of an embedded body on a timescale much shorter than the disk’s viscous time. This mode of migration is commonly referred to as Type I, while a gap-opening planet co-evolving with its disk is said to undergo Type II migration Ward (1997). Hahn & Ward (1996) suggested that the potentially large differences between Type I and Type II timescales could lead to bodies of the former class being delivered to, and then resonantly trapped by, bodies of the latter class. The concentration of a significant mass of

¹IAU Working Group on Extrasolar planets, <http://www.ciw.edu/boss/IAU/div3/wgesp/planets.shtml>; California and Carnegie Planet Search, <http://www.exoplanets.org>

protoplanets in exterior MMRs of a pre-existing giant planet could, they argued, promote the formation of subsequent giant planets. In this work, we undertake the first part of a numerical investigation of this scenario. We make use of two different approaches. In the first and simpler one, we take a standard N-body code and add to it forces to mimic the effects of planet-disk interaction, guided by analytic work as well as by numerical simulations. The second approach is to take this same N-body code and add to it a simple disk model, which viscously co-evolves with embedded planets and interacts with them. The disk model is one-dimensional; that is, it describes an azimuthally-averaged thin disk, with disk-planet torques obtained from analytic formulae rather than from a full calculation of the launching of density waves by the planet. This approach is intermediate between the above “N-body with disk forces” and an actual 2- (or 3-) dimensional hydrodynamic simulation. With both approaches, we find that for typical protoplanetary disk parameters, single Type I bodies have a high likelihood of being resonantly captured by Type II bodies. We further demonstrate that significant numbers of protoplanets can occupy adjacent resonances, or indeed share a single resonance, of a giant planet. This paper is organized as follows: In §2, we give a brief summary of the theory of planet-disk interactions. In §3.1, we present simulations performed using the simpler of our two approaches, while in §3.2 we repeat these simulations using our hybrid “N-body+viscous disk” code. We discuss our results in §4, and summarize them in §5.

2. Planet-disk torques, migration and eccentricity evolution

Here, we provide a brief summary of the theory of planet-disk interaction, focusing on the results we will make use of in this paper. A body orbiting in a gas disk launches density waves at Lindblad resonances, exchanging energy and angular momentum with the disk; the disk thus modifies the body’s orbit. The net effect on the body can be obtained by summing over all resonances (Goldreich & Tremaine 1980; see also e.g. Goldreich & Sari 2003). One begins by expanding the gravitational potential of the perturbing body in a double Fourier series. The l, m component has a pattern speed

$$\Omega_{l,m} = \Omega_p + (l - m)\kappa_p/m \quad (1)$$

where Ω_p and κ_p are the azimuthal and epicyclic frequencies, respectively, of the planet. There is a Lindblad resonance between this potential component and disk material at a radius r in the disk where

$$\Omega(r) \pm \frac{\kappa(r)}{m} = \Omega_{l,m}. \quad (2)$$

Also, a corotation resonance occurs where

$$\Omega(r) = \Omega_{l,m} \quad (3)$$

The radial pressure gradient in the disk causes both the epicyclic and the azimuthal frequency to differ slightly from their Keplerian values:

$$\Omega^2(r) = \frac{GM_*}{r^3} + \frac{1}{r\rho} \frac{\partial}{\partial r}(\rho c_s^2) \quad (4)$$

and

$$\kappa^2(r) = \frac{1}{r^3} \frac{\partial}{\partial r} (r^4 \Omega^2). \quad (5)$$

In the case of a planet on a circular orbit, all components with $l \neq m$ are zero; $\Omega_{m,m} = \Omega_p$, so all components have the pattern speed of the planet. Furthermore, there is no torque contribution at all from corotation resonances. We will focus on this simpler situation for the moment. In summing the effect of all Lindblad resonances, one can define a torque density, or torque per unit disk radius. Following Ward (1997), the torque density experienced by the disk due to a planet of mass $M = \mu M_*$ on a circular orbit about a primary of mass M_* at radius r_p , is

$$\left[\frac{dT}{dr}(r) \right]_{LR} = \text{sgn}(r - r_p) \frac{2\mu^2 \Sigma(r) r_p^4 \Omega_p^4}{r(1 + 4\xi^2) \kappa^2} m^4 \psi^2 \quad (6)$$

where $\xi \equiv m c_s / r \kappa$, c_s is the gas sound speed, and Ψ is the dimensionless satellite forcing function,

$$\Psi = \frac{\pi}{2} \left[\frac{1}{m} \left| \frac{db_{1/2}^m}{d\beta} \right| + 2 \frac{\Omega}{\kappa} \sqrt{1 + \xi^2} \beta_{1/2}^m(\beta) \right], \quad (7)$$

with $\beta \equiv r/r_p$ and

$$b_{1/2}^m(\beta) = \frac{2}{\pi} \int_0^\pi \frac{\cos m\theta d\theta}{\sqrt{1 - 2\beta \cos \theta + \beta^2}} \quad (8)$$

being the Laplace coefficient. Following Menou & Goodman (2004), we account in a simple, approximate way for the finite thickness of the disk by replacing the Laplace coefficient with a softened approximation,

$$b_{1/2}^m(\beta) \approx \frac{2}{\pi \beta^{1/2}} K_0 \left(m \sqrt{\beta - 2 + \frac{1}{\beta} + \frac{(BH)^2}{rr_p}} \right) \quad (9)$$

with K_0 the modified Bessel function of the second kind, order 0, $H \approx c_s/\Omega$ the disk scale height, and B a dimensionless scaling factor. In going from a summing of torques at discrete

resonances to a torque density, the wavenumber m is turned into a continuous function of radius:

$$m(r) = \left[\frac{\kappa^2}{(\Omega - \Omega_p)^2 - c_s^2/r^2} \right]^{1/2}. \quad (10)$$

Ward (1988), Artymowicz (1993) and Papaloizou & Larwood (2000) study planet-disk torques from an eccentric planet. In particular, Papaloizou & Larwood (2000) look at the case where eccentricity exceeds disk aspect ratio H/r . They sum over enough $l \neq m$ torques for convergence, in order to obtain the net effect on the planet’s orbit. However, they neglect corotation torques, as these are zero for their model surface density profile ($\Sigma \propto r^{-3/2}$). They obtain approximate fits for the timescales of angular momentum and eccentricity change:

$$t_m \equiv -\frac{J}{\dot{J}} = 3.5 \times 10^5 f_s^{1.75} \left[\frac{1 + \left(\frac{e_p a_p}{1.3H}\right)^5}{1 - \left(\frac{e_p r_p}{1.1H}\right)^4} \right] \left(\frac{H/a_p}{0.07} \right)^2 \left(\frac{\Sigma}{588 \text{gcm}^{-2}} \right)^{-1} \left(\frac{M_p}{1M_\oplus} \right)^{-1} \left(\frac{a_p}{1AU} \right)^{-1/2} \text{yrs} \quad (11)$$

and

$$t_e \equiv -\frac{e}{\dot{e}} = 2.5 \times 10^3 f_s^{2.5} \left[1 + \frac{1}{4} \left(\frac{e_p}{H/a_p} \right)^3 \right] \left(\frac{H/a_p}{0.07} \right)^2 \left(\frac{\Sigma}{588 \text{gcm}^{-2}} \right)^{-1} \left(\frac{M_p}{1M_\oplus} \right)^{-1} \left(\frac{a_p}{1AU} \right)^{-1/2} \text{yrs} \quad (12)$$

In the limit of low planetary eccentricities, these results are broadly consistent with the earlier work of Ward (1997) and Artymowicz (1993). Most noteworthy is the short orbital decay timescale; for constant eccentricity, $a/\dot{a} = t_m/2$, so from Eq. 11, it is of order 10^4 years for a body of mass $10 M_\oplus$ in a typical model disk. This would seem to present a significant problem for the core-accretion model of giant planet formation, since the assembly of the solid core likely takes a million years or more (e.g. Lissauer 1987, Kokubo & Ida 2000, Thommes, Duncan, & Levison 2003, Inaba, Wetherill, & Ikoma 2003), with perhaps several million more required to reach runaway gas accretion (itself a very fast process) and acquire a Jovian atmosphere (Pollack et al. 1996).

If a planet does manage to grow to something approaching gas giant mass before plunging into its parent star, the problem of rapid inward migration becomes less severe. Torque increases with planet mass; for a sufficiently massive body, $\sim 10^2 M_\oplus$ for typical disk models, the torques overcome disk viscosity and an annular gap about the orbit is opened in the gas (Goldreich & Tremaine 1980), provided that the Hill (Roche) radius also exceeds the disk scale height (Lin & Papaloizou 1986, 1993). A planet opening a clean gap prevents disk material from crossing its orbit, and thus becomes locked into the disk’s viscous evolution. In general, this entails a sharp decrease in the rate of orbital decay.

The eccentricity evolution of a gap-opening planet is as yet not well understood. In the gapless case, the dominant effect on the planet’s eccentricity is damping by coorbital

Lindblad resonances. When a gap forms and empties or strongly depletes the coorbital region, the evolution of the eccentricity comes down to a delicate balance between the effect of non-coorbital eccentric Lindblad resonances (driving) and corotation resonances (damping). Though the latter are slightly stronger than the former, Goldreich & Sari (2003) and Ogilvie & Lubow (2003) argue that corotation resonances in fact saturate, thus favouring net eccentricity excitation. However, Goldreich & Sari (2003) suggest that damping may be reestablished once a planet’s eccentricity becomes so high that its radial excursion exceeds the gap width. In contrast, numerical simulations tend to show only damping of eccentricities (e.g. Kley, Peitz, & Bryden 2004) unless the planet mass approaches that of a brown dwarf (Papaloizou, Nelson, & Masset 2001).

3. Numerical simulations

For the purpose of the subsequent numerical experiments we will assume that, the above difficulties notwithstanding, a protostellar disk does manage to form one initial gas giant planet, massive enough to open a gap and orbitally co-evolve with its parent disk. Taking this as our initial condition, we then study the evolution of sub-gap-opening bodies originating at larger radii than the gas giant. To better assess the robustness of our results, we repeat the investigation with two different numerical approaches.

3.1. N-body + dissipative disk forces

Simulations are performed with a code based on SyMBA (Symplectic Massive Body Algorithm), an N-body program optimized for near-Keplerian systems (Duncan, Levison, & Lee 1998). SyMBA is based on the symplectic orbit-integration algorithm of Wisdom & Holman (1991), with its high efficiency and good energy-conservation properties, and adds an adaptive timestep in order to resolve close encounters between bodies. Using a model gas disk, we then calculate the angular momentum and eccentricity decay rate for each sub-gap-opening body from Eqs. 11 and 12 each timestep, and apply the corresponding additional acceleration $\vec{a} = a_\phi \hat{\phi} + a_r \hat{r}$, where

$$a_\phi = -\frac{\dot{J}}{M_p r_p} = \frac{1}{t_m} \frac{J}{M_p r_p} \quad (13)$$

and

$$a_r = -2 \frac{\vec{v} \cdot \vec{r}}{r^2 t_e}. \quad (14)$$

The giant planet in our simulations is treated differently, however. For simplicity, we prevent it from migrating altogether. This is done in a way that mimics, rather simplistically, the effect of being locked in a disk gap: We choose a radius r_{gap} for the center of the gap and a fractional gap “width” w , then apply an azimuthal acceleration a'_ϕ as follows:

$$a'_\phi = \frac{a - r_{\text{gap}}}{|a - r_{\text{gap}}| + wa} a_\phi \quad (15)$$

Since a_ϕ is negative, a'_ϕ is negative at $a > r_{\text{gap}}$, positive at $a < r_{\text{gap}}$, and falls off smoothly to zero as $a \rightarrow r_{\text{gap}}$. Therefore, if our “gap-opening” planet is displaced from the gap midpoint, it experiences a restoring torque towards $a = r_{\text{gap}}$.

3.1.1. Single growing protoplanet

We begin with a simple case: A $300 M_\oplus$ (approximately Jupiter-mass) body starts on a circular orbit at 5 AU about a $1 M_\odot$ star, centered in a “disk gap” as described above. We place a second body of mass $1 M_\oplus$ on a circular orbit at 15 AU. A model disk with surface density

$$\Sigma_g = 1000 \left(\frac{r}{1\text{AU}} \right)^{-1} \text{ g cm}^{-2} \quad (16)$$

and scale height

$$H = 0.03 \left(\frac{r}{1\text{AU}} \right)^{5/4} r \quad (17)$$

is used to calculate dJ/dt and de/dt . After 5×10^5 years of simulation time, the outer body’s mass is increased linearly in time to $30 M_\oplus$ over the next million years. The resulting orbital evolution is shown in Fig. 1. At its original mass of $1 M_\oplus$, the outer body migrates inward until it is captured in the larger body’s 1:2 exterior mean motion resonance. Once its mass has reached $M \approx 2.5 M_\oplus$, at $t \approx 5.5$ Myrs, it jumps from the 1:2 to the 2:3 MMR. It remains there through the rest of its growth and until the end of the simulation at 5 Myrs, showing no signs of orbital instability. We perform other simulations with different migration rates; we achieve this by simply scaling the disk surface density uniformly, while keeping the disk power law index and scale height profile the same (note that $J/\dot{J} \propto H^2 \Sigma_g^{-1}$ for small eccentricity). Some of these other results are also shown in Fig. 1. When $\Sigma(1\text{AU}) = 1500 \text{ g cm}^{-2}$, the growing body undergoes a second resonance jump, from the 2:3 to the 3:4 MMR. When $\Sigma(1\text{AU}) = 6000 \text{ g cm}^{-2}$, the body does the same, then also leaves the 3:4 resonance, but no closer resonance is able to capture it, and it gets all the way to the gap-opener’s orbit (and in the case pictured, physically collides with it). Our experiment thus suggests that, for parameters relevant to a protoplanetary disk— $\Sigma_g \sim (10^2 - 10^3) \text{ g cm}^{-2}$, $H/r \sim (10^{-2} - 10^{-1})$ —single bodies of gas giant core/ice giant mass, $\sim (10^0 - 10^1) M_\oplus$, have

a high likelihood of being captured upon encountering a first-order (n:n+1) mean-motion resonance of a gas giant-sized body.

3.1.2. Multiple protoplanets

We now turn to the question of what happens when multiple sub-gap-opening bodies encounter a gas giant. Dropping several fully-formed Neptune-mass planets into a disk makes for a rather questionable initial condition; as in the previous section, we therefore begin with small bodies whose masses we increase over time. This time, we do so in a manner which, at least in a broad qualitative sense, conforms to how planet formation proceeds. We make use of the prescription for post-runaway or “oligarchic” (Kokubo & Ida 1998) growth in the form given by Thommes, Duncan, & Levison (2003). This describes the mass accretion rate of a mass M_{pp} protoplanet growing together with an oligarchy of similar neighbours, all embedded in a disk of much smaller planetesimals of characteristic mass m_{plsm} :

$$\frac{dM_{pp}}{dt} \approx AM_{pp}^{2/3} (\Sigma_s - BM^{2/3}), \quad (18)$$

where Σ_s is the initial local surface density of the planetesimal disk, and A and B are given by

$$A = \frac{3.9b^{2/5}C_D^{2/5}G^{1/2}M_*^{1/6}\rho_g^{2/5}}{\rho_{plsm}^{4/15}\rho_{pp}^{1/3}a_{pp}^{1/10}m_{plsm}^{2/15}}, \quad B = \frac{0.23M_*^{1/3}}{ba_{pp}^2} \quad (19)$$

C_D is the drag coefficient, a dimensionless constant ≈ 1 ; b is the spacing in Hill radii between adjacent protoplanets; M_* is the mass of the central star; $\rho_g \approx \Sigma_g/H$ is the (volume) density of gas; and ρ_{pp} and ρ_{plsm} are the densities of protoplanet and planetesimals, respectively.

We put protoplanets, initially of mass $0.1 M_\oplus$, on circular orbits between 8 and 20 AU. Their masses are changed over time according to Eq. 18, with Σ_s multiplied by a randomly-generated factor between 0.25 and 1.75 in order to mimic the stochastic nature of the accretion. Σ_g and H are given by Eqs. 16 and 17. We use $b = 10$, $\rho_{pp} = \rho_{plsm} = 1.5 \text{ g cm}^{-3}$, $m_{plsm} = 10^{-18} M_\oplus$; the latter two amount to a planetesimal size of $\sim 0.01 \text{ km}$.

A typical example of the orbital evolution which results is shown in Fig. 2. The outer bodies accelerate their inward migration as they grow in mass, and one after another is resonantly captured by the “Jupiter” at 5 AU. As they continue to grow and the applied torque increases, some bodies jump to a closer-in resonance, analogous to the behavior seen in Fig. 1. At the end of the simulation at 5 Myrs, five bodies share the 2:3 MMR with the gas giant. A sixth body is further outward; examination of the different possible resonance angles (e.g. Murray & Dermott 1999) shows it to be in a 4:7 MMR with the gas giant, and simultaneously, in a 6:7 MMR with the bodies which are themselves in the 3:2 resonance with the




Fig. 1.— Simulation of growing, sub-gap-opening (Type I) body in a gas disk, using the “N-body + dissipative disk forces” scheme. The simulated disk has a surface density of $\Sigma_{1\text{AU}}(r/\text{AU})^{-1} \text{ g cm}^{-2}$. In the top panel, the orbital evolution is shown for three cases: $\Sigma_{1\text{AU}} = 1000 \text{ g cm}^{-2}$ (black), $\Sigma_{1\text{AU}} = 1500 \text{ g cm}^{-2}$ (green), and $\Sigma_{1\text{AU}} = 6000 \text{ g cm}^{-2}$ (red). In each case the body’s semimajor axis a , pericenter $q \equiv a(1 - e)$ and apocenter $Q \equiv a(1 + e)$ are plotted. A body of mass $300 M_{\oplus}$ orbits at 5 AU; the Type I body spends time trapped in several of its mean-motion resonances (blue dashed lines). The time evolution of the Type I body’s mass is the same in each of the three cases, and is shown in the bottom panel.

giant ($4 : 7 = 2 : 3 \times 6 : 7$). Meanwhile, all the 2:3 bodies are in a 1:1 commensurability with each other. This seemingly precarious configuration is attainable here because the tendency of the bodies to gravitationally scatter each other as they approach to such close range, is kept in check by the strong damping of eccentricities which is simultaneously applied. Other outcomes are shown in Fig. 3. The 2:3 resonance is strongly favoured as the final location of the growing protoplanets, with the smallest bodies tending to end up further out, in a 5:9 (5:6) or 4:7 (6:7) MMR with the gas giant (the 2:3 MMR bodies), respectively.

3.2. Hybrid N-body + viscous disk

We now repeat our experiments with a different numerical approach. The code used here is called VDisk-SyMBA; as the name suggests, it also uses SyMBA for evolving planetary bodies, but the disk is treated in a more sophisticated way: We co-evolve with the planets a one-dimensional disk model, which not only changes in time due to its own viscosity, but also exchanges angular momentum with the N-body part, modifying its own density profile as well as the orbits of the planets. A similar computational approach, though without the N-body component, was utilized by Lin & Papaloizou (1986) to follow the evolution of single protoplanets in a disk. Since radius is its only dimension, the disk is of necessity vertically and azimuthally averaged. However, both vertical and azimuthal structure are *implicitly* included in the model. For the former, we simply assign a scale height; for the latter, planet-disk interaction—a process which after all works by raising azimuthally asymmetric perturbations in the disk—is handled by way of the torque density formulation of Eq. 6 applied to the azimuthally averaged disk surface density. This approach to simulating planet-disk interactions is thus intermediate in complexity between our first method and a full hydrodynamic simulation. The fidelity of the planet-disk interactions is only as good as the torque density used, but because the disk is one-dimensional, this code has a significant speed advantage over two- or three-dimensional hydrodynamic codes. In practice we find that millions of years can typically be simulated in one day of real time on a desktop-class machine, thus simulations spanning the entire lifetime of a protoplanetary gas disk, constrained observationally to be $\sim 10^7$ years or less (Haisch, Lada, & Lada 2001) are within easy reach.

The gas disk is divided into radial bins of arbitrary size at the start of the simulation. The most stringent upper limit on bin size comes from the necessity of properly resolving planet-disk torques; accurately tracking the disk’s viscous self-evolution is much more forgiving in that respect. We generally find it necessary to scale the bin size to be some fraction of the local disk scale height.




Fig. 2.— A simulation of multiple growing protoplanets outside a giant planet (fixed at 5 AU) using the “N-body + dissipative disk forces” scheme. Top panels: Orbital evolution of semimajor axis, peri- and apocenters of the bodies. The left panel shows the initial migration, while the one on the right shows the full simulation with a smaller vertical scale. Bottom left: Evolution of the protoplanet masses by the “oligarchy” prescription. Bottom right: Final state showing resonance locations.




Fig. 3.— Snapshots taken at 5×10^6 years of a set of ten simulations (differing only in the initial orbital phases of the bodies) of a population of growing protoplanets on orbits exterior to a gap-opening gas giant. The locations of some of the giant’s exterior mean-motion resonance centers are shown (vertical lines). Most of the surviving protoplanets are locked in the 2:3 resonance, typically with one additional body in the 4:7 and 5:9 resonance. In one case (G), there is a single much larger protoplanet, formed from the merger of several smaller ones, in the 3:4 resonance.

The governing equation for the surface density of a Keplerian viscous disk is

$$\frac{\partial \Sigma}{\partial t} = \frac{1}{r} \frac{\partial}{\partial r} \left[3r^{1/2} \frac{\partial}{\partial r} (\nu \Sigma r^{1/2}) - \frac{r^{1/2}}{\pi \sqrt{GM_*}} \frac{\partial T}{\partial r} \right] \quad (20)$$

where $\partial T / \partial r$ is the torque density in the disk due to the planet (Eq. 6). For the viscous self-evolution of the disk, we solve this equation each timestep, with the second term set to zero, using an explicit upwind differencing scheme. The planet-disk torques are computed and applied to both planet and disk in a separate step. An important part of this is to compute Ω in each bin; although we assume perfectly Keplerian gas velocities for the disk’s viscous evolution, the torque asymmetry felt by sub-gap-opening bodies is, as mentioned in §2, primarily due to the sub-Keplerian gas orbital speed shifting resonance locations. We calculate Ω in each bin at each timestep using the finite-difference version of Eq. 4. We then make the approximation that $\kappa = \Omega$.

The planet-disk torques are obtained using the expression for torque density, Eq. 6. We are thus computing the torques for a circular orbit; when $e \neq 0$, we use a simplified version of Eq. 12 (without the eccentricity-dependent part in square brackets) to separately change the eccentricity. For gas surface density Σ_g , we average among the value at the semimajor axis, pericentre and apocentre of the orbit. This is a somewhat ad-hoc prescription for the eccentricity evolution, but it does reproduce the analytically-obtained results for a non-gap-opening body, while for a gap-opening planet, damping of eccentricities does not commence until the planet’s radial excursion causes it to overrun its gap. Thus we have implemented a compromise of sorts between numerical results on the one hand (e.g. Papaloizou, Nelson, & Masset 2001; Kley, Peitz, & Bryden 2004) and the analytic work of Goldreich & Sari (2003) and Ogilvie & Lubow (2003) on the other. It should also be pointed out that implicit in our approach is the assumption that angular momentum is deposited in the disk right at the location of a resonance. This is a good assumption as long as the torquing planet has r_H greater than the scale height H of the disk, in this case waves launched at resonances will indeed be nonlinear and shock immediately, thus depositing their angular momentum right away (Lin & Papaloizou 1993). However, for bodies having $r_H < H$, a more accurate treatment would include modelling the nonlocal deposition of angular momentum in the disk (Takeuchi, Miyama, & Lin 1996; Rafikov 2002). The effect of our simplification is that we underestimate the gap-opening threshold mass for bodies with $r_H < H$.

3.2.1. *Single protoplanet*

We generate an initial gas disk of similar surface density and scale height profile as in §3.1.1. We adopt a standard α -parameterization for the viscosity of the disk, $\nu = \alpha c_s H$,

with $\alpha = 10^{-4}$. A $300 M_{\oplus}$ body is, again, started on a circular orbit at 5 AU, while a $1 M_{\oplus}$ body is started on a circular orbit at 15 AU. As before, after 5×10^5 years, its mass is linearly increased to $30 M_{\oplus}$ over the next 10^6 years. The orbital evolution of both bodies, together with the disk, is shown in Fig. 4

The most obvious difference in the evolution, with respect to Fig. 1, is that the inner planet also migrates. It immediately opens a gap in the disk, initially moving outward slightly as the nearby disk material rearranges itself about the freshly-opened gap. Thereafter, the planet moves inward along with the accretion flow of the disk, going from 5 AU to 2 AU in just under 2 Myrs. The outer planet migrates much faster, at a similar rate as for the $\Sigma_g(1\text{AU}) = 1000 \text{ g cm}^{-2}$ case in §3.1.1, and is captured into the 1:2 MMR of the inner planet after a bit more than 10^5 years. Unlike the cases shown in Fig. 1, it remains in the 1:2 resonance as it grows. The strength with which the outer body is pushed toward the inner body is somewhat reduced by the inward migration of the inner body, as well as by the time evolution of the gas disk (Σ_g decreases as the disk spreads). Also, as can be clearly seen in Fig. 4, as the outer body becomes more massive (from $\sim 8 \times 10^5$ years onward) it begins to open a gap of its own. As this happens, gas begins to be trapped between the two bodies; the more effective the trapping, the more this gas acts to push the bodies apart, counteracting the asymmetric torque felt by the outer body. Another difference with respect to Fig. 1 is that the gap-opening body temporarily acquires a substantial eccentricity, such that its radial excursion becomes equal to the gap width, where the onset of strong damping then halts eccentricity growth. At the same time, the smaller body’s eccentricity decreases temporarily (examination of the 2:1 resonance angles shows a coincident change in their libration centers). This behavior appears to arise in a fairly narrow range of parameter space, wherein the torque on the outer body is, for an extended period of time, just below the threshold needed to move it from the 1:2 to 2:3 resonance. It is not apparent in any of the other simulations presented below, and we defer a closer examination to future work.

We perform a second simulation with a more massive gas disk, $\Sigma_g \approx 3000(r/1 \text{ AU})^{-1} \text{ g cm}^{-2}$. The results are shown in Fig. 5; this time, the outer planet goes from the 1:2 to the 2:3 resonance with the inner almost as soon as its growth commences. This puts it right at the outer edge of the gap.

3.3. Multiple protoplanets

Using the same prescription as in §3.1.2, we again grow multiple protoplanets originating as $0.1 M_{\oplus}$ bodies exterior to the gas giant. A typical simulation is shown in Fig. 6. In addition, snapshots at 1.5×10^6 years of a set of similar simulations (differing only in the




Fig. 4.— Orbital evolution of an inner gap-opening (Type II) $300 M_{\oplus}$ planet and an outer sub-gap-opening (Type I), initially $1 M_{\oplus}$ body. For each body, the semimajor axis a (black), pericenter $q \equiv a(1 - e)$ and apocenter $Q \equiv a(1 + e)$ (both green) are plotted. The locations of the gap-opening body's 1:2 and 2:3 mean-motion resonances (MMRs) are also shown (upper and lower yellow dashed line, respectively). The $1 M_{\oplus}$ body grows linearly to $30 M_{\oplus}$ between 5×10^5 yrs and 1.5×10^6 yrs; it remains trapped in the 1:2 MMR until the end of




Fig. 5.— The simulation shown in Fig. 4, repeated with a higher initial gas disk surface density of $\Sigma \approx 3000(r/1 \text{ AU})^{-1} \text{ g cm}^{-2}$. This time, the outer body transitions from the 1:2 to the 2:3 MMR at 5×10^5 years.

randomly-generated initial orbital phases of the bodies) are shown in Fig 7. All surviving protoplanets at this time are locked in either the 2:3 or the 1:2 exterior MMR of the gap-opening giant, with the majority residing in the former. As in the simulations of §3.1.2, multiple protoplanets—as many as four in this case—end up sharing the 2:3 resonance.

4. Discussion

The numerical simulations we have performed here suggest that when rapidly-migrating, sub-gap-opening protoplanets approach a gap-opening gas giant, resonant capture is a likely outcome. The two different approaches we use in our simulations—first, dissipative forces added to an N-body code, and then, a more self-consistent treatment of the disk which includes gap formation and viscous accretion—both produce similar results. A particularly interesting feature common to both sets of simulation results is the “stacking” of multiple multi-Earth-mass bodies in a single mean-motion resonance of the gas giant. Such seemingly precarious resonant configurations can be assembled in our simulations because the eccentricity damping we apply to sub-gap bodies stabilizes them against catastrophic mutual scattering with pre-existing resonant bodies as they approach a MMR.

The ability of a giant planet locked into the viscous transport of the disk to arrest the orbital decay of $O(10^0 - 10^1) M_{\oplus}$ bodies, which, at least in a simple disk model like the one used here, move inward much faster than the disk material, has important implications for giant planet formation. In the classic core-accretion model (Pollack et al. 1996), most of a giant planet’s “childhood”, several Myrs in length, is spent as a $\sim 10^1 M_{\oplus}$ body waiting to accrete a massive gas envelope. However, the work of Ward (1997) and e.g. Papaloizou & Larwood (2000), summarized in §2, implies an orbital decay timescale for such bodies of only $\sim 10^4$ years. Thus, having cores held up by a more slowly-migrating gas giant can potentially be a big help in enabling them to reach core mass at all. However, even a gap-opening planet does not offer a safe haven indefinitely. As can be seen in the simulations of §3.2, the gas giant undergoes significant migration in only 2 Myrs, moving in more than 1 AU/Myr on average. And this, it should be stressed, is with a rather low α viscosity parameter of 10^{-4} . With $\alpha = 10^{-3}$, a gap-opening planet starting at 5 AU has well below 1 Myr before it plunges into the star, barring some sort of “parking mechanism” at the inner edge of the disk (e.g. a central magnetospheric cavity; Lin, Bodenheimer, & Richardson 1996). In other words, for a sufficiently large disk viscosity, Type II migration becomes just as problematically fast as Type I migration. Thus, in order for a gas giant to be of use as a safety net for smaller bodies, it must exist in a disk, or a region of a disk, that has fairly low viscosity. Just such a possibility is offered by the layered accretion or “dead




Fig. 6.— Simulation of the orbital evolution of a population of growing protoplanets outside a gap-opening gas giant, plotted as in Figs. 4 and 5. The protoplanets migrate inward rapidly, and one protoplanet after another encounters the exterior mean-motion resonances of the giant. After a period of vigorous interactions lasting until $\sim 8 \times 10^5$ years, the protoplanets all occupy either the 1:2 or the 2:3 MMR.




Fig. 7.— Snapshots taken at 1.5×10^6 years of a set of ten simulations (differing only in the initial orbital phases of the bodies) of a population of growing protoplanets on orbits exterior to a gap-opening gas giant. The locations of the exterior 2:3 and 1:2 mean-motion resonances of the gap-opening planet are shown as vertical lines. All surviving protoplanets are locked in one or the other resonance.

zone” disk model (Gammie 1996; Matsumura & Pudritz 2003), in which the bulk of the disk transports angular momentum only at small radii, $\lesssim 1$ AU, and beyond several AU. In between, angular momentum is only transported in the disk’s surface layers. This is because the assumed source of disk viscosity, magnetorotational instability (Balbus & Hawley 1991), only operates in a sufficiently ionized disk; at small radii the star can provide this ionization and at larger radii cosmic rays take over, but at intermediate radii, the column depth of the disk is too high to allow cosmic rays to penetrate all the way through. The result is that from less than 1 AU to several AU (the exact bounds depend on the details of the model) the disk has a low viscosity, and thus gas—and gap-embedded planets—will accrete slowly through this region, giving planet formation more time to play out. Even then, the ultimate survival of whatever planets are produced may require the removal of at least part of the outer disk by photoevaporation rather than accretion (Shu, Johnstone, & Hollenbach 1993; Hollenbach, Yorke, & Johnstone 2000; Matsuyama, Johnstone, & Murray 2003). It has been suggested that a giant planet may promote the accretion of further planets due to the enhancement of disk surface density near the edges of a freshly-excavated gap (Lin & Papaloizou 1979). The outer gap edge, in particular, may locally concentrate solids because the transition from super- to sub-Keplerian gas orbital speed there ought to cause a pile-up of planetesimals in a narrow radial range (Bryden et al. 2000). We demonstrate that, in agreement with the suggestion of Hahn & Ward (1996), an analogous effect exists when solids grow to larger size, where disk tides rather than aerodynamic drag are the primary agent of gas-solids interaction. Indeed, both effects—the pile-up of small planetesimals and the resonant capture of large protoplanets—are likely to operate in concert. This brings up a shortcoming in our prescription for protoplanet growth: We use an estimate based on orderly post-runaway (“oligarchic”) growth, which assumes a smooth surface density of solids and a uniform spacing of neighbouring protoplanets. Both assumptions will certainly not hold up well when a local enhancement of solids exists, and especially when multiple bodies are competing to accrete in the *same* mean-motion resonance. However, our intent here has been merely to make the accretion prescription plausible in the broadest sense; a more detailed study of the accretion of multiple resonantly-locked, co-orbital protoplanets is a subject for future work.

A shortcoming common to both of our simulation schemes is that we neglect the effect of corotation resonances on sub-gap-opening planets. Since their strength is proportional to the gradient of the quantity $(\Omega\Sigma_g/\kappa^2)$, this is more of an issue for gap-opening planets, which have large Σ_g gradients at the gap edges. These resonances are pivotal in determining the rate—indeed the sign—of the eccentricity evolution of planets in gaps (Goldreich & Sari 2003; Ogilvie & Lubow 2003). In modeling gap-opening planets, we also make no explicit use of corotation resonance torques, but rather use a compromise among the competing models

for the net effect of all resonances on the eccentricity evolution: Eccentricity is neither driven nor damped while the planet is fully in its gap, but is damped once it becomes eccentric enough to run into the gap edges. A more general caveat is that our simulation results are contingent on the validity of treating planet-disk torques of multiple nearby bodies in an azimuthally-averaged, linearly-superposed manner. It will be very interesting to see to what extent full hydrodynamic simulations reproduce the resonant configurations we have generated here, especially the extreme case of multiple Type I bodies sharing a Type II body’s MMR.

The simulations performed here lend support to the idea that the pre-existence of a giant planet makes the formation of subsequent planets easier. If so, the formation of the first giant planet may be the critical event in a protoplanetary disk. But with no “safety net” of its own, how can the first solid core be assembled fast enough? Menou & Goodman (2004) offer one possibility: In a more complex disk model than is usually used for such calculations (including the ones in this work), opacity transitions can create local migration bottlenecks which stretch the Type I timescale to $\sim 10^6$ years. The work of Cuzzi & Zahnle (2004) is also suggestive: They show that icy planetesimals undergoing orbital decay due to aerodynamic gas drag can produce and maintain a massive plume of water vapor once they reach the water sublimation radius (“snow line”). If much of this water recondenses just outside the snow line, a large local enhancement in the surface density of solids may result. At that radius, accretion will be more rapid and produce bigger bodies than elsewhere. Furthermore, although we have been working from the perspective of the core accretion model, one can also imagine a hybrid scenario wherein the first giant planet forms rapidly via the competing disk instability model (Cameron 1978, Boss 1997, Mayer et al. 2002), thus avoiding the Type I issue altogether, and subsequently facilitates the formation of its successors by core accretion. This initial, innermost planet (or planets) may well end up migrating into the star, leaving only the successors to populate the mature system. That could circumvent some of the inconsistencies with observations commonly associated with the disk instability model, namely higher-than-detected planet masses and (in the context of our Solar System) insufficiently enhanced planet metallicities. Finally, regardless of the formation mechanism, it is worth remembering that only about a quarter of detected multi-planet systems contain resonant pairs. Therefore, an important question to be addressed in future work is how readily mean-motion resonances can be broken as a system evolves.

5. Summary

We have performed numerical simulations which suggest that gap-opening (Type II) gas giant-sized bodies are very efficient at resonantly capturing sub-gap-opening (Type I) bodies of Earth to Neptune size in their exterior mean-motion resonances. In fact, multiple such bodies may accumulate in a single resonance. The local concentration of such a large amount of mass for significant lengths of time stands in sharp contrast to the problem which arises when considering the core-accretion growth of a single giant planet in isolation, namely that the growth timescale exceeds the migration timescale long before the critical mass for runaway accretion of a gas envelope ($\sim 10^1 M_{\oplus}$) is reached. Hence, the *first* gas giant to form may be the most important, paving the way for its successors. Also, these results suggest one way to build mean-motion resonant systems of giant planets like those discovered by radial-velocity surveys to be orbiting GJ 876, HD 82943, and 55 Cnc: Rather than giant planets forming separately and then undergoing convergent migration, resonance locking may happen much earlier, when the second planet is still just a growing core.

I would like to thank G. Bryden, W. Kley, D. Lin, M.-H. Lee, G. Marcy and F. Masset for stimulating and informative discussions, the Kavli Institute for Theoretical Physics for its hospitality during the early part of this work, and an anonymous referee for valuable comments and suggestions. This work is supported by the Natural Sciences and Engineering Research Council of Canada.

REFERENCES

- Artymowicz, P. 1993, ApJ, 419, 166
- Balbus, S. A., & Hawley, J. F. 1991, ApJ, 376, 214
- Boss, A. P. 1997, Science, 276, 1836
- Bryden, G., Różyczka, M., Lin, D. N. C., & Bodenheimer, P. 2000, ApJ, 540, 1091
- Cameron, A. G. W. 1978, Moon and Planets, 18, 5
- Cuzzi, J. N., & Zahnle, K. J. 2004, ApJ, 614, 490
- Duncan, M. J., Levison, H. F., & Lee, M. H. 1998, AJ, 116, 2067
- Gammie, C. F. 1996, ApJ, 457, 355

- Goldreich, P., & Sari, R. 2003, *ApJ*, 585, 1024
- Goldreich, P., & Tremaine, S. 1980, *ApJ*, 241, 425
- Hahn, J. M., & Ward, W. R. 1996, in *Lunar and Planetary Institute Conference Abstracts*, 479–+
- Haisch, K. E., Lada, E. A., & Lada, C. J. 2001, *ApJ*, 553, L153
- Hollenbach, D. J., Yorke, H. W., & Johnstone, D. 2000, *Protostars and Planets IV*, 401
- Inaba, S., Wetherill, G. W., & Ikoma, M. 2003, *Icarus*, 166, 46
- Kley, W. 2000, *MNRAS*, 313, L47
- Kley, W., Peitz, J., & Bryden, G. 2004, *A&A*, 414, 735
- Kokubo, E., & Ida, S. 1998, *Icarus*, 131, 171
- . 2000, *Icarus*, 143, 15
- Lee, M. H., & Peale, S. J. 2002, *ApJ*, 567, 596
- Lin, D. N. C., Bodenheimer, P., & Richardson, D. C. 1996, *Nature*, 380, 606
- Lin, D. N. C., & Papaloizou, J. 1979, *MNRAS*, 188, 191
- . 1986, *ApJ*, 309, 846
- Lin, D. N. C., & Papaloizou, J. C. B. 1993, in *Protostars and Planets III*, 749–835
- Lissauer, J. J. 1987, *Icarus*, 69, 249
- Masset, F., & Snellgrove, M. 2001, *MNRAS*, 320, L55
- Matsumura, S., & Pudritz, R. E. 2003, *ApJ*, 598, 645
- Matsuyama, I., Johnstone, D., & Murray, N. 2003, *ApJ*, 585, L143
- Mayer, L., Quinn, T., Wadsley, J., & Stadel, J. 2002, *Science*, 298, 1756
- Menou, K., & Goodman, J. 2004, *ApJ*, 606, 520
- Murray, C. D., & Dermott, S. F. 1999, *Solar System Dynamics* (Cambridge University Press)
- Murray, N., Paskowitz, M., & Holman, M. 2002, *ApJ*, 565, 608

- Nelson, R. P., & Papaloizou, J. C. B. 2002, MNRAS, 333, L26
- Ogilvie, G. I., & Lubow, S. H. 2003, ApJ, 587, 398
- Papaloizou, J. C. B., & Larwood, J. D. 2000, MNRAS, 315, 823
- Papaloizou, J. C. B., Nelson, R. P., & Masset, F. 2001, A&A, 366, 263
- Pollack, J. B., Hubickyj, O., Bodenheimer, P., Lissauer, J. J., Podolak, M., & Greenzweig, Y. 1996, Icarus, 124, 62
- Rafikov, R. R. 2002, ApJ, 572, 566
- Shu, F. H., Johnstone, D., & Hollenbach, D. 1993, Icarus, 106, 92
- Takeuchi, T., Miyama, S. M., & Lin, D. N. C. 1996, ApJ, 460, 832
- Thommes, E. W., Duncan, M. J., & Levison, H. F. 2003, Icarus, 161, 431
- Ward, W. R. 1988, Icarus, 73, 330
- . 1997, Icarus, 126, 261
- Wisdom, J., & Holman, M. 1991, AJ, 102, 1528